



**Part - B (Chemistry)**

51. Given below are half cell reaction:



$$E_{\text{Mn}^{2+}/\text{MnO}_4^-}^\circ = 1.1510 \text{ V}$$



$$E_{\text{O}_2/\text{H}_2\text{O}}^\circ = +1.223 \text{ V}$$

Will the permanganate ion, MnO_4^- liberate O_2 from water in the presence of an acid?

(1) Yes, because $E_{\text{cell}}^\circ = +2.733 \text{ V}$

(2) No, because $E_{\text{cell}}^\circ = -2.733 \text{ V}$

(3) Yes, because $E_{\text{cell}}^\circ = +0.287 \text{ V}$

(4) No, because $E_{\text{cell}}^\circ = -0.287 \text{ V}$

Ans: [3]

52. Identify the incorrect statement from the following

(1) Ionisation enthalpy of alkali metals decreases from top to bottom in the group

(2) Lithium is the strongest reducing agent among the alkali metals

(3) Alkali metals react with water to form their hydroxides

(4) The oxidation number of K in KO_2 is +4

Ans: [4]

53. Identify the incorrect statement from the following

(1) In an atom, all the five 3d orbitals are equal in energy in free state

(2) The shapes of d_{xy} , d_{yz} , and d_{zx} orbitals are similar to each other and $d_{x^2-y^2}$ and d_{z^2} are similar to each other

(3) All the five 5d orbitals are different in size when compared to the respective 4d orbitals

(4) All the five 4d orbitals have shapes similar to the respective 3d orbitals

Ans: [2]

54. Gadolinium has a low value of third ionisation enthalpy because of

(1) High electronegativity (2) High basic character

(3) Small size (4) High exchange enthalpy

Ans: [4]

55. The IUPAC name of an element with atomic number 119 is

(1) ununium (2) ununochium

(3) ununennium (4) unnilennium

Ans: [3]

56. Match List I with List II

List I

List II

(Hydrides)

(Nature)

(a) MgH_2

(i) Electron deficient

(b) GeH_4

(ii) Electron rich

(c) B_2H_6

(iii) Electron deficient

(d) HF


(iv) Ionic

Choose the correct answer from the options given below.

(1) a - i, b - ii, c - iv, d - iii (2) a - ii, b - iii, c - iv, d - i

(3) a - iv, b - i, c - ii, d - iii (4) a - iii, b - i, c - ii, d - iv

Ans: [3]



What is Y in the above reaction?

(1) $\text{RCOO}^- \text{X}^+$

(2) $(\text{RCOO})_2\text{Mg}$

(3) $\text{RCOO}^- \text{Mg}^+ \text{X}$

(4) $\text{R}_3\text{CO}^- \text{Mg}^+ \text{X}$

Ans: [3]

58. Match List I with List II

List I

List II

(Drug class)

(Drug molecule)

(a) Antacids

(i) Salvarsan

(b) Antihistamines

(ii) Morphine

(c) Analgesics

(iii) Cimetidine

(d) Antimicrobials

(iv) Seldane

Choose the correct answer from the options given below:

(1) a - i, b - iv, c - ii, d - iii (2) a - iv, b - iii, c - i, d - ii

(3) a - iii, b - ii, c - iv, d - i (4) a - iii, b - iv, c - ii, d - i

Ans: [4]

59. Which of the following statement is not correct about diborane

- (1) The four terminal Hydrogen atoms and the two Boron atoms lie in one plane
- (2) Both the Boron atoms are sp^2 hybridised
- (3) There are two 3-centre-2-electron bonds
- (4) The four terminal B-H bonds are two centre two electron bonds

Ans: [2]

60. The incorrect statement regarding enzymes is :

- (1) Enzymes are polysaccharides
- (2) Enzymes are very specific for a particular reaction and substrate
- (3) Enzymes are biocatalysts
- (4) Like chemical catalysts enzymes reduce the activation energy of bio processes

Ans: [1]

61. Given below are two statements : one is labelled Assertion (A) and the other is labelled as Reason (R).

Assertion (A) : In a particular point defect, an ionic solid electrically neutral even if few of its cations missing from its unit cells.


Reason (R) : In an ionic solid, Frenkel defect arises due to dislocation of cation from its lattice site to interstitial site, maintaining overall electrical neutrality.

In the light of the above statements, choose the appropriate answer from the options given below

- (1) (A) is correct but (R) is not correct
- (2) (A) is not correct but (R) is correct
- (3) Both (A) and (R) are correct and (R) is correct explanation of (A)
- (4) Both (A) and (R) are correct but (R) is not correct explanation of (A)

Ans: [4]

62. Which of the following P-V curve represents maximum work done?



Ans: [4]

63. Given below are two statements:

Statement I : Primary aliphatic amines react with HNO_2 to give unstable diazonium salts

Statement II : Primary aromatic amines react with HNO_2 to form diazonium salts which are stable even above 300 K.

In the light of the above statements, choose the most appropriate answer from the options given below:

- (1) Statement I is correct but Statement II is incorrect
- (2) Statement I is incorrect but Statement II is correct
- (3) Both Statement I and Statement II are correct
- (4) Both Statement I and Statement II are incorrect

Ans: [1]


64. The IUPAC name of the complex -



- (1) dicyanidosilver(I) diaquaargentate(I)
- (2) diaquasilver(I) dicyanidoargentate(I)
- (3) dicyanidosilver(II) diaquaargentate(II)
- (4) diaquasilver(II) dicyanidoargentate(II)

Ans: [2]


65. Which compound amongst the following is not in aromatic compound?



Ans: [2]

66. Given below are two statements:

Statement I : In the coagulation of a negative sol, the flocculating power of the three given ions in the order-



Statement II : In the coagulation of a positive sol, the flocculating power of the three given salts in the order



In the light of the above statements, choose the most appropriate answer from the options given below :

- (1) Statement I is correct but Statement II is incorrect
 (2) Statement I is incorrect but Statement II is correct
 (3) Both Statement I and Statement II are correct
 (4) Both Statement I and Statement II are incorrect

Ans: [1]

67. Match List I with List II

List I	List II
(a) Li	(i) absorbent for carbon dioxide
(b) Na	(ii) electrochemical cells
(c) KOH	(iii) coolant in fast breeder reactions
(d) Cs	(iv) photoelectric cell

Choose the correct answer from the options given below

- (1) a - i, b - iii, c - iv, d - ii (2) a - ii, b - iii, c - i, d - iv
 (3) a - iv, b - i, c - iii, d - ii (4) a - iii, b - iv, c - ii, d - i

Ans: [2]

68. Given below are two statement : one is labelled as Assertion (A) and the other is labelled as Reason (R).

Assertion (A) : ICl is more reactive than I_2 .

Reason (R) : I - Cl bond is weaker than I - I bond.

In the light of the above statements, choose the most appropriate answer from the options given below:

- (1) (A) is correct but (R) is correct
 (2) (A) is not correct but (R) is correct
 (3) Both (A) and (R) are correct and (R) is the correct explanation of (A)
 (4) Both (A) and (R) are correct but (R) is not the correct explanation of (A)

Ans: [3]

69. Amongst the following which one will have maximum lone pair - lone pair' electron repulsions?

- (1) SF_4 (2) XeF_2
 (3) ClF_3 (4) IF_5

Ans: [2]

70. Given below are two statements:

Statement I : The acidic strength of monosubstituted nitrophenol is higher than phenol because of electron withdrawing nitro group.

Statement II : o-nitrophenol, m-nitrophenol and p-nitrophenol will have same acidic strength as they have one nitro group attached to the phenolic ring.

In the light of the above statements, choose the most appropriate answer from the options given below:

- (1) Statement I is correct but Statement II is incorrect
 (2) Statement I is incorrect but Statement I is correct
 (3) Both Statement I and Statement II are correct
 (4) Both Statement I and Statement II are incorrect

Ans: [1]

71. Which one is not correct mathematical equation for Dalton's Law of partial pressure? Here p = total pressure of gaseous mixture.

- (1) $P_i = \chi_i P$, where P_i = partial pressure of i^{th} gas
 χ_i = mole fraction of i^{th} gas in gaseous mixture
 (2) $P_i = \chi_i P_i^\circ$ where χ_i = mole fraction of i^{th} gas in gaseous mixture

P_i° = pressure of i^{th} gas in pure state

- (3) $P = P_1 + P_2 + P_3$

- (4) $P = n_1 \frac{RT}{V} + n_2 \frac{RT}{V} + n_3 \frac{RT}{V}$

Ans: [2]

72. Given below are two statements:

Statement I : The boiling points the following hydrides of group 16 elements increases in the order :



Statement II : The boiling points of these hydrides increase with increase in molar mass.

In the light of the above statements choose the most appropriate answer from the options given below.

- (1) Statement I is correct but Statement II is incorrect
 (2) Statement I is incorrect but Statement II is correct
 (3) Both Statement I and Statement II are correct
 (4) Both Statement I and Statement II are incorrect

Ans: [2]

73. In one molal solution that contains 0.5 mole of a solute, there is

- (1) 100 mL of solvent (2) 1000 g of solvent
 (3) 500 mL of solvent (4) 500 g of solvent


Ans: [4]

74. Choose the correct statement

- (1) Diamond is sp^3 hybridised and graphite is sp^2 hybridized
 (2) Both diamond and graphite are used as dry lubricants
 (3) Diamond and graphite have two dimensional network
 (4) Diamond is covalent and graphite is ionic

Ans: [1]

75. What mass of 95% pure CaCO_3 will be required to neutralise 50 M HCl solution according to the following reaction?





[Calculate upto second place of decimal point]

- (1) 3.65 g (2) 9.50 g
 (3) 1.25 g (4) 1.32 g

Ans: [4]

76. Which of the following is suitable to synthesize chlorobenzene?



- (3) Benzene, Cl_2 , anhydrous FeCl_3
 (4) Phenol, NaNO_2 , HCl, CuCl

Ans: [3]

77. The pH of the solution containing 50 mL each of 0.10 M sodium acetate and 0.01 M acetic acid is.

[Given pK_a of $\text{CH}_3\text{COOH} = 4.57$]

- (1) 4.57 (2) 2.57
 (3) 5.57 (4) 3.57

Ans: [3]

78. Which amongst the following is incorrect statement?

- (1) H_2^+ ion has one electron
 (2) O_2^+ ion is diamagnetic
 (3) The bond orders of O_2^+ , O_2 , O_2^- and O_2^{2-} are 2.5, 2, 1.5 and 1, respectively
 (4) C_2 molecule has four electrons in its two degenerate π molecular orbitals

Ans: [2]

79. Which statement regarding polymers is not correct?

- (1) Thermoplastic polymers are capable of repeatedly softening and hardening on heating and cooling respectively
 (2) Thermosetting polymers are reusable
 (3) Elastomers have polymer chains held together by weak intermolecular forces
 (4) Fibers possess high tensile strength

Ans: [2]

80. Match List I and List II

List I	List II
(Procucts formed)	(Reaction of carbonyl)
(a) Cyanohydrin	(i) NH_2OH
(b) Acetal	(ii) RNH_2
(c) Schiff's base	(iii) alcohol
(d) Oxime	(iv) HCN

Choose the correct answer from the option given below:

- (1) a - i, b - iii, c - ii, d - iv (2) a - iv, b - iii, c - ii, d - i
 (3) a - iii, b - iv, c - ii, d - i (4) a - ii, b - iii, c - iv, d - i


Ans: [2]
81. Given below are two statements:

Statement I : The boiling points of aldehydes and ketones are higher than hydrocarbons of comparable molecular masses because of weak molecular association in aldehydes and ketones due to dipole - dipole interactions.

Statement II : The boiling points of aldehydes and ketones are lower than the alcohols of similar molecular masses due to the absence of H-bonding

In the light of the above statement, choose the most appropriate answer from the options given below.

- (1) Statement I is correct but Statement II is incorrect
 (2) Statement I is incorrect but Statement II is correct
 (3) Both Statement I and Statement II are correct
 (4) Both Statement I and Statement II are incorrect

Ans: [3]
82. The given graph is a representation of kinetics of a reaction


The y and x axes for zero and first order reactions, respectively are :

- (1) zero order (y = rate and x = concentration)
 first order (y = $t_{1/2}$ and x = concentration)
 (2) zero order (y = rate and x = concentration)
 first order (y = rate and x = $t_{1/2}$)
 (3) zero order (y = concentration and x = time)


first order (y = $t_{1/2}$ and x = concentration)

(4) zero order (y = concentration and x = time)

first order (y = rate constant and x = concentration)

Ans: [1]
83. The incorrect statement regarding chirality is:

- (1) Enantiomers are superimposable mirror images on each other
 (2) A racemic mixture shows zero optical rotation
 (3) $\text{S}_{\text{N}}1$ reaction yields 1 : 1 mixture of both enantiomers
 (4) The product obtained by $\text{S}_{\text{N}}2$ reaction of haloalkane having chirality at the reactive site shows inversion of configuration

Ans: [1]
84. The Kjeldahl's method for the estimation of nitrogen can be used to estimate the amount of nitrogen in which one of the following compounds?

Ans: [1]
85. At 298 K, the standard electrode potentials of $\text{Cu}^{2+} / \text{Cu}$, $\text{Zn}^{2+} / \text{Zn}$, $\text{Fe}^{2+} / \text{Fe}$ and $\text{Ag}^{2+} / \text{Ag}$ are 0.34 V, -0.76 V, -0.44 V and 0.80 V, respectively,

On the basis of standard electrode potential, predict which of the following reaction cannot occur?

- (1) $\text{FeSO}_4(\text{aq}) + \text{Zn}(\text{s}) \rightarrow \text{ZnSO}_4(\text{aq}) + \text{Fe}(\text{s})$
 (2) $2\text{CuSO}_4(\text{aq}) + 2\text{Ag}(\text{s}) \rightarrow 2\text{Cu}(\text{s}) + \text{Ag}_2\text{SO}_4(\text{aq})$
 (3) $\text{CuSO}_4(\text{aq}) + \text{Zn}(\text{s}) \rightarrow \text{ZnSO}_4(\text{aq}) + \text{Cu}(\text{s})$
 (4) $\text{CuSO}_4(\text{aq}) + \text{Fe}(\text{s}) \rightarrow \text{FeSO}_4(\text{aq}) + \text{Cu}(\text{s})$

Ans: [2]

86. Match List I with List II





List I	List II
(Ores)	(Composition)
(a) Haematite	(i) Fe_3O_4
(b) Magnetite	(ii) ZnCO_3
(c) Calamine	(iii) Fe_2O_3
(d) Kaolinite	(iv) $[\text{Al}_2(\text{OH})_4\text{Si}_2\text{O}_5]$

Choose the correct answer from the options given below:

- (1) a - iii, b - i, c - iv, d - ii (2) a - i, b - iii, c - ii, d - iv
 (3) a - i, b - ii, c - iii, d - iv (4) a - iii, b - i, c - ii, d - iv

Ans: [4]

87. Which one of the following is not formed when acetone reacts with 2-pentanone in the presence of dilute NaOH followed by heating?



Ans: [4]

88. Given below are two statements

Statement I : In Lucas test, primary, secondary and tertiary alcohols are distinguished on the basis of their reactivity with conc. $\text{HCl} + \text{ZnCl}_2$, known as Lucas Reagent.

Statement II : Primary alcohols are most reactive and immediately produce turbidity at room temperature on reaction with Lucas Reagent

In the light of the above statements, choose the most appropriate answer from the options given below:

- (1) Statement I is correct but Statement II is incorrect
 (2) Statement I is incorrect but Statement II is correct
 (3) Both Statement I and Statement II are correct
 (4) Both Statement I and Statement II are incorrect

Ans: [1]

89. If radius of second Bohr orbit of the He^+ ion is 105.8 pm, what is the radius of third Bohr orbit of Li^{2+} ion?

- (1) 1.587 pm (2) 1.58.7 Å
 (3) 1.58.7 pm (4) 15.87 pm

Ans: [3]


90. A 10.0 L flask contains 64 g of oxygen at 27°C (Assume O_2 gas is behaving ideally). The pressure inside the flask in bar is

(Given $R = 0.0831 \text{ L bar K}^{-1} \text{ mol}^{-1}$)

- (1) 49.8 (2) 4.9
 (3) 2.5 (4) 498.6

Ans: [2]

91. Find the emf of the cell in which the following reaction takes place at 298 K




(Given that $E_{\text{cell}}^\circ = 10.5 \text{ V}$, $\frac{0.303 RT}{F} = 0.059$ at 298 K)

- (1) 0.9615 V (2) 1.05 V
 (3) 1.0385 V (4) 1.385 V

Ans: [Bonus]

92. The correct IUPAC name of the following compounds is



- (1) 1-bromo-4-methyl-5-chlorohexane-3-ol
 (2) 6-bromo-4-methyl-2-chlorohexane-4-ol
 (3) 1-bromo-5-methyl-4-chlorohexane-3-ol
 (4) 6-bromo-2-methyl-4-chlorohexane-4-ol

Ans: [3]

93. In the neutral or faintly alkaline medium KMnO_4 oxidises iodide into iodate. The change in oxidation state of manganese in this reaction is from

- (1) +7 to +3 (2) +6 to +5
 (3) +7 to +4 (4) +6 to +4

Ans: [3]

94. $3\text{O}_2(\text{g}) \rightleftharpoons 2\text{O}_3(\text{g})$

for the above reaction at 298 K, K_c is found to be 3.0×10^{-59} . If the concentration of O_2 at equilibrium is 0.040 M then concentration of O_3 in M is

- (1) 2.4×10^{31} (2) 1.2×10^{21}
 (3) 4.38×10^{-32} (4) 1.9×10^{-63}

Ans: [3]

95. Copper crystallises in fcc unit cell with cell edge length of 3.608×10^{-8} cm. The density of copper is 8.92 g dm^{-3} . Calculate the atomic mass of copper

- (1) 60 u (2) 65 u
 (3) 63.1 u (4) 31.55 u

Ans: [3]

96. The pollution due to oxides of sulphur gets enhanced due to the presence of :



- (a) particulate matter
 (b) ozone
 (c) hydrocarbons
 (d) hydrogen peroxide

Choose the most appropriate answer from the options given below:

- (1) (b), (c), (d) only (2) (a), (c), (d) only
 (3) (a), (d) only (4) (a), (b), (d) only

Ans: [4]

97. The product formed from the following reaction sequence is



Ans: [2]

98. Compound X on reaction with O_3 followed by $\text{Zn} / \text{H}_2\text{O}$ gives formaldehyde and 2-methyl propanal as products. The compound X is

- (1) 2-Methylbut-2-ene (2) Pent-2-ene
 (3) 3-Methylbut-1-ene (4) 2-Methylbut-1-ene

Ans: [3]

99. For a first order reaction $\text{A} \rightarrow \text{Products}$, initial concentration of A is 0.1 M, which becomes 0.001 M after 5 minutes. Rate constant for the reaction in min^{-1} is

- (1) 0.4606 (2) 0.2303
 (3) 1.3818 (4) 0.9212

Ans: [4]

100. The order of energy absorbed which is responsible for the color of complex

- (A) $[\text{Ni}(\text{H}_2\text{O})_2(\text{en})_2]^{2+}$
 (B) $[\text{Ni}(\text{H}_2\text{O})_4(\text{en})]^{2+}$ and
 (C) $[\text{Ni}(\text{en})_3]^{2+}$ is

- (1) (C) > (A) > (B) (2) (B) > (A) > (C)
 (3) (A) > (B) > (C) (4) (C) > (B) > (A)

Ans: [1]